

Rad17 (Phospho Ser635) Rabbit pAb

CatalogNo: YP1457

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA, IHC

MW

- 77kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho peptide around human Rad17 (Ser635)

Specificity This antibody detects endogenous levels of Human Rad17 (phospho-Ser635)

Target Information

Gene name RAD17 R24L

Protein Name Rad17 (Ser635)


Organism	Gene ID	UniProt ID
Human	5884 ;	O75943 ;
Mouse	19356 ;	Q6NXW6 ;

Cellular Localization Nucleus . Phosphorylated form redistributes to discrete nuclear foci upon DNA damage.

Tissue specificity Overexpressed in various cancer cell lines and in colon carcinoma (at protein level) . Isoform 2 and isoform 3 are the most abundant isoforms in non irradiated cells (at protein level) . Ubiquitous at low levels. Highly expressed in testis , where it is expressed within the germinal epithelium of the seminiferous tubuli. Weakly expressed in seminomas (testicular tumors) .

Function Function:Essential for sustained cell growth , maintenance of chromosomal stability , and ATR-dependent checkpoint activation upon DNA damage. Has a weak ATPase activity required for binding to chromatin. Participates in the recruitment of the RAD1-RAD9-HUS1 complex onto chromatin , and in CHEK1 activation. May also serve as a sensor of DNA replication progression , and may be involved in homologous recombination. ,induction:By X-ray irradiation (isoform 1 , isoform 3 and isoform 4) . ,PTM:Phosphorylated. Phosphorylation on Ser-646 and Ser-656 is cell cycle-regulated , enhanced by genotoxic stress , and required for activation of checkpoint signaling. Phosphorylation is mediated by ATR upon UV or replication arrest , whereas it may be mediated both by ATR and ATM upon ionizing radiation. Phosphorylation on both sites is required for interaction with RAD1 but dispensable for interaction with RFC3 or RFC4. ,similarity:Belongs to the rad17/RAD24 family. ,subcellular location:Phosphorylated form redistributes to discrete nuclear foci upon DNA damage. ,subunit:Part of a DNA-binding complex containing RFC2 , RFC3 , RFC4 and RFC5. Interacts with RAD1 and RAD9 within the RAD1-RAD9-HUS1 complex. Interacts with RAD9B , POLE , NHP2L1 and MCM7. DNA damage promotes interaction with ATR or ATM and disrupts interaction with the RAD1-RAD9-HUS1 complex. ,tissue specificity:Overexpressed in various cancer cell lines and in colon carcinoma (at protein level) . Isoform 2 and isoform 3 are the most abundant isoforms in non irradiated cells (at protein level) . Ubiquitous at low levels. Highly expressed in testis , where it is expressed within the germinal epithelium of the seminiferous tubuli. Weakly expressed in seminomas (testicular tumors) . ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: http://www.immunoway.com.cn
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**Rad17 (Phospho
Ser635) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)