

SGK3 (Phospho Thr320) Rabbit pAb

CatalogNo: YP1486

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC

MW

- 54kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho peptide around human SGK3 (Thr320)

Specificity This antibody detects endogenous levels of Human SGK3 (phospho-Thr320)

Target Information

Gene name SGK3 CISK SGKL

Protein Name Serine/threonine-protein kinase Sgk3

Organism	Gene ID	UniProt ID
Human	100533105 ;	Q96BR1 ;
Mouse	170755 ;	Q9ERE3 ;
Rat		Q8R4V0 ;

Cellular Localization

Cytoplasmic vesicle . Early endosome. Recycling endosome. Endosomal localization is a prerequisite for complete kinase activity. It is essential for its colocalization with the kinase responsible for phosphorylating Ser-486 thus allowing PDPK1 phosphorylation of Thr-320 resulting in complete activation of SGK3. Localized in vesicle-like structures and in the early endosome. Colocalizes with SLC9A3/NHE3 in the recycling endosomes.


Tissue specificity

Expressed in most tissues with highest levels in pancreas , kidney liver , heart and brain and lower levels in lung , placenta and skeletal muscle. Expression is higher in ER-positive breast tumors than ER-negative breast tumors.

Function

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,enzyme regulation:Two specific sites , one in the kinase domain (Thr-320) and the other in the C-terminal regulatory region (Ser-486) , need to be phosphorylated for its full activation. ,Function:Involved in the activation of potassium channels. Mediates cell IL-3-dependent survival signals. ,similarity:Belongs to the protein kinase superfamily. ,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. ,similarity:Contains 1 AGC-kinase C-terminal domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 PX (phox homology) domain. ,subcellular location:Localized in vesicle-like structures and in the early endosome. ,tissue specificity:Expressed in most tissues with highest levels in pancreas , kidney liver , heart and brain and lower levels in lung , placenta and skeletal muscle. ,

| Validation Data



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SGK3 (Phospho Thr320) Rabbit pAb

