

## NCoA-3 (Phospho Thr24) Rabbit pAb

CatalogNo: YP1510

### | Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 160kD (Observed)

#### Isotype

- IgG

### | Recommended Dilution Ratios

WB 1:1000-2000

### | Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### | Basic Information

**Clonality** Polyclonal

### | Immunogen Information

**Immunogen** Synthesized phospho peptide around human SRC-3 (Thr24)

**Specificity** This antibody detects endogenous levels of Human SRC-3 (phospho-Thr24)

### | Target Information

**Gene name** NCOA3 AIB1 BHLHE42 RAC3 TRAM1

<b>Protein Name</b>	SRC-3 (Thr24)		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">8202;</a>	<a href="#">Q9Y6Q9;</a>
	Mouse		<a href="#">O09000;</a>
	Rat		<a href="#">Q9EPU2;</a>
<b>Cellular Localization</b>	Cytoplasm. Nucleus. Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus.		
<b>Tissue specificity</b>	Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.		
<b>Function</b>	<p>Alternative products:Additional isoforms seem to exist,Catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,Domain:Contains three Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. Motifs 1 and 2 are essential for the association with nuclear receptors, and constitute the RID domain (Receptor-interacting domain).,enzyme regulation:Coactivator activity on nuclear receptors and NF-kappa-B pathways is enhanced by various hormones, and the TNF cytokine, respectively. TNF stimulation probably enhances phosphorylation, which in turn activates coactivator function. In contrast, acetylation by CREBBP apparently suppresses coactivation of target genes by disrupting its association with nuclear receptors.,Function:Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit.,miscellaneous:NCOA3 is frequently amplified or overexpressed in breast and ovarian cancers.,polymorphism:The length of the poly-Gln region is polymorphic in the normal population.,PTM:Acetylated by CREBBP. Acetylation occurs in the RID domain, and disrupts the interaction with nuclear receptors and regulates its function.,PTM:Methylated by CARM1.,PTM:Phosphorylated by IKK complex. Regulated its function.,similarity:Belongs to the SRC/p160 nuclear receptor coactivator family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 PAS (PER-ARNT-SIM) domain.,subcellular location:Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus.,subunit:Interacts with CARM1 (By similarity). Present in a complex containing NCOA2, IKKA, IKKB, IKBKG and the histone acetyltransferase protein CREBBP. Interacts with CASP8AP2, NR3C1 and PCAF. Interacts with ATAD2 and this interaction is enhanced by estradiol.,tissue specificity:Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.,</p>		

| Validation Data

| Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**NCoA-3 (Phospho  
Thr24) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)