

NFAT1 (Phospho Ser326) Rabbit pAb

CatalogNo: YP1563 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA, IHC

MW

- 100kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human NFAT1 (Phospho Ser326)

Specificity This antibody detects endogenous levels of Human, Mouse, Rat NFAT1 (Phospho Ser326). The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): KTSPD

| Target Information

Gene name NFATC2 NFAT1 NFATP

Protein Name NFAT1 (Phospho Ser326)


Organism	Gene ID	UniProt ID
Human	4773 ;	Q13469 ;
Mouse	18019 ;	Q60591 ;

Cellular Localization Cytoplasm. Nucleus. Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription.


Tissue specificity Expressed in thymus , spleen , heart , testis , brain , placenta , muscle and pancreas. Isoform 1 is highly expressed in the small intestine , heart , testis , prostate , thymus , placenta and thyroid. Isoform 3 is highly expressed in stomach , uterus , placenta , trachea and thyroid.

Function Alternative products:Additional isoforms seem to exist ,Domain:Rel Similarity Domain (RSD) allows DNA-binding and cooperative interactions with AP1 factors. ,Function:Plays a role in the inducible expression of cytokine genes in T-cells , especially in the induction of the IL-2 , IL-3 , IL-4 , TNF-alpha or GM-CSF. ,induction:Inducibly expressed in T-lymphocytes upon activation of the T-cell receptor (TCR) complex. Induced after co-addition of phorbol 12-myristate 13-acetate (PMA) and ionomycin. ,PTM:In resting cells , phosphorylated by NFATC-kinase on at least 18 sites in the 99-363 region. Upon cell stimulation , all these sites except Ser-243 are dephosphorylated by calcineurin. Dephosphorylation induces a conformational change that simultaneously exposes an NLS and masks an NES , which results in nuclear localization. Simultaneously , Ser-53 or Ser-56 is phosphorylated; which is required for full transcriptional activity. ,similarity:Contains 1 RHD (Rel-like) domain. ,subcellular location:Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription. ,subunit:Member of the multicomponent NFATC transcription complex that consists of at least two components , a pre-existing cytoplasmic component NFATC2 and an inducible nuclear component NFATC1. Other members such as NFATC4 , NFATC3 or members of the activating protein-1 family , MAF , GATA4 and Cbp/p300 can also bind the complex. The phosphorylated form specifically interacts with XPO1; which mediates nuclear export. NFATC proteins bind to DNA as monomers. Interacts with NFATC2IP. ,tissue specificity:Expressed in thymus , spleen , heart , testis , brain , placenta , muscle and pancreas. ,

| Validation Data



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).



Western Blot analysis of 1 MCF-7 treated with LPS, 2 MCF-7, using primary antibody at 1:1000 dilution. Secondary antibody (catalog#:RS23920) was diluted at 1:10000

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
NFAT1 (Phospho Ser326) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)