

PDK1 (Phospho Thr338) Rabbit pAb

CatalogNo: YP1634

Key Features

Host Species

- Rabbit

Reactivity

- Human,Mouse,Rat

Applications

- WB,ELISA,IHC

MW

- 50kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human PDHK1 (Phospho Thr338)

Specificity This antibody detects endogenous levels of Human,Mouse,Rat PDHK1 (Phospho Thr338).The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):T(p)A

Target Information

Gene name PDK1 PDHK1

Protein Name PDHK1 (Phospho Thr338)


Organism	Gene ID	UniProt ID
Human	5163 ;	Q15118 ;
Mouse	228026 ;	Q8BFP9 ;
Rat		Q63065 ;

Cellular Localization Mitochondrion matrix .

Tissue specificity Expressed predominantly in the heart. Detected at lower levels in liver , skeletal muscle and pancreas.

Function Catalytic activity:ATP + [pyruvate dehydrogenase (acetyl-transferring)] = ADP + [pyruvate dehydrogenase (acetyl-transferring)] phosphate. ,Function:Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit , thus contributing to the regulation of glucose metabolism. ,similarity:Belongs to the PDK/BCKDK protein kinase family. ,similarity:Contains 1 histidine kinase domain. ,tissue specificity:Expressed predominantly in the heart. ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
PDK1 (Phospho Thr338) Rabbit pAb

