

CDK4 (Phospho Thr172) Rabbit pAb

CatalogNo: YP1663

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 33kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human CDK4 (Phospho-Thr172)

Specificity This antibody detects endogenous levels of CDK4 (Phospho-Thr172) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):ALtPV

Target Information

Gene name CDK4
Protein Name CDK4 (Phospho-Thr172)

Organism	Gene ID	UniProt ID
Human	1019;	P11802;
Mouse	12567;	P30285;
Rat	94201;	P35426;

Cellular Localization Cytoplasm . Nucleus . Nucleus membrane . Cytoplasmic when non-complexed. Forms a cyclin D-CDK4 complex in the cytoplasm as cells progress through G (1) phase. The complex accumulates on the nuclear membrane and enters the nucleus on transition from G (1) to S phase. Also present in nucleoli and heterochromatin lumps. Colocalizes with RB1 after release into the nucleus. .

Tissue specificity Brain ,Muscle ,

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Disease:CDK4 mutations are involved in tumor formation. ,Disease:Defects in CDK4 are the cause of cutaneous malignant melanoma 3 (CMM3) [MIM:609048 , 155600]. Malignant melanoma is a malignant neoplasm of melanocytes , arising de novo or from a preexisting benign nevus , which occurs most often in the skin but also may involve other sites. ,enzyme regulation:Phosphorylation at Thr-172 is necessary for enzymatic activity. ,Function:Probably involved in the control of the cell cycle. ,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily. ,similarity:Contains 1 protein kinase domain. ,subunit:Forms a stable complex with D-type G1 cyclins. Interacts with SEI1 and ZNF655/VIK. ,

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
CDK4 (Phospho Thr172) Rabbit pAb