

## eNOS (Phospho Ser633) Rabbit pAb

CatalogNo: YP1704

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 132kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

WB 1:500-2000

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from human eNOS (Phospho-Ser633)

**Specificity** This antibody detects endogenous levels of eNOS (Phospho-Ser633) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):KEsSN

---

## | Target Information

**Gene name** NOS3

**Protein Name** eNOS (Phospho-Ser633)

Organism	Gene ID	UniProt ID
Human	<a href="#">4846</a> ;	<a href="#">P29474</a> ;
Mouse	<a href="#">18127</a> ;	<a href="#">P70313</a> ;
Rat	<a href="#">24600</a> ;	<a href="#">Q62600</a> ;


**Cellular Localization** Cell membrane. Membrane , caveola. Cytoplasm , cytoskeleton. Golgi apparatus. Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity.

**Tissue specificity** Platelets , placenta , liver and kidney.

**Function** Catalytic activity:L-arginine + n NADPH + n H (+) + m O (2) = citrulline + nitric oxide + n NADP (+) . ,cofactor: Binds 1 FAD. ,cofactor: Binds 1 FMN. ,cofactor: Heme group. ,cofactor: Tetrahydrobiopterin (BH4) . May stabilize the dimeric form of the enzyme. ,enzyme regulation: Stimulated by calcium/calmodulin. Inhibited by NOSIP and NOSTRIN. ,Function: Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF) -induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets. ,online information: Nitric oxide synthase entry ,polymorphism: Variation in NOS3 seem to be associated with susceptibility to coronary spasm. ,similarity: Belongs to the NOS family. ,similarity: Contains 1 FAD-binding FR-type domain. ,similarity: Contains 1 flavodoxin-like domain. ,subcellular location: Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity. ,subunit: Homodimer. Interacts with NOSIP and NOSTRIN. ,tissue specificity: Platelets , placenta , liver and kidney. ,

---

## | Validation Data



Western Blot analysis of mouse heart ,using primary antibody at 1:1000 dilution. Secondary antibody (catalog#:RS23920) was diluted at 1:10000

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**eNOS (Phospho Ser633) Rabbit pAb**