

## TOPBP1 (Phospho Ser1159) Rabbit pAb

CatalogNo: YP1741 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 167kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

WB 1:500-2000

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from human TOPBP1 (Phospho-Ser1159)

**Specificity** This antibody detects endogenous levels of TOPBP1 (Phospho-Ser1159) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): LAsNL

---

## | Target Information

**Gene name** TOPBP1 KIAA0259

**Protein Name** TOPBP1 (Phospho-Ser1159)

Organism	Gene ID	UniProt ID
Human	<a href="#">11073</a> ;	<a href="#">Q92547</a> ;
Mouse	<a href="#">235559</a> ;	<a href="#">Q6ZQF0</a> ;

**Cellular Localization**

Nucleus. Cytoplasm , cytoskeleton , microtubule organizing center , centrosome. Cytoplasm , cytoskeleton , spindle pole. Chromosome. Detected on unpaired autosomes in meiotic prophase cells. Detected on X and Y chromosomes during later stages of prophase. Colocalizes with ATR and H2AX at unsynapsed chromosome cores during prophase (By similarity) . Has a uniform nuclear distribution during G phase. Colocalizes with BRCA1 at stalled replication forks during S phase. In mitotic cells it colocalizes with BRCA1 at spindle poles and centrosomes during metaphase and anaphase. Detected in discrete foci together with PML and numerous DNA repair enzymes after DNA damage by alkylating agents , UV or gamma irradiation. Localizes to sites of DNA damage in a H2AX- independent manner. .


**Tissue specificity** Highly expressed in heart , brain , placenta , lung and kidney.

**Function**

Function:Required for DNA replication. Plays a role in the rescue of stalled replication forks and checkpoint control. Binds double-stranded DNA breaks and nicks as well as single-stranded DNA. Recruits the SWI/SNF chromatin remodeling complex to E2F1-responsive promoters. Down-regulates E2F1 activity and inhibits E2F1-dependent apoptosis during G1/S transition and after DNA damage. Induces a large increase in the kinase activity of ATR. ,induction:Up-regulated during the S phase of the cell cycle. Up-regulated by E2F1 and interferon. ,PTM:Phosphorylated on serine and threonine residues in response to X-ray irradiation. ,PTM:Ubiquitinated and degraded by the proteasome. X-ray irradiation reduces ubiquitination. ,similarity:Contains 8 BRCT domains. ,subcellular location:Detected on unpaired autosomes in meiotic prophase cells. Detected on X and Y chromosomes during later stages of prophase. Colocalizes with ATR and H2AFX at unsynapsed chromosome cores during prophase (By similarity) . Has a uniform nuclear distribution during G phase. Colocalizes with BRCA1 at stalled replication forks during S phase. In mitotic cells it colocalizes with BRCA1 at spindle poles and centrosomes during metaphase and anaphase. Detected in discrete foci together with PML and numerous DNA repair enzymes after DNA damage by alkylating agents , UV or gamma irradiation. ,subunit:Binds E2F1 , PML , RAD9A , SMARCA2 , SMARCA4 , POLE and UBR5. May interact with TOP2B. ,tissue specificity:Highly expressed in heart , brain , placenta , lung and kidney. ,

---

## | Validation Data



Western Blot analysis of 1HeLa cell 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody (catalog#:RS23920) was diluted at 1:10000

## Contact information

Orders: order.cn@immunoway.com  
 Support: support.cn@immunoway.com  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**TOPBP1 (Phospho Ser1159) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)