

RSK2 (Phospho Tyr529) Rabbit pAb

CatalogNo: YP1775

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 81kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human RSK2 (Phospho-Tyr529)

Specificity This antibody detects endogenous levels of RSK2 (Phospho-Tyr529) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): VEyLH

| Target Information

Gene name RPS6KA3 ISPK1 MAPKAPK1B RSK2

Protein Name RSK2 (Phospho-Tyr529)


Organism	Gene ID	UniProt ID
Human	6197 ;	P51812 ;
Mouse	110651 ;	P18654 ;

Cellular Localization Nucleus . Cytoplasm .

Tissue specificity Expressed in many tissues , highest levels in skeletal muscle.

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,cofactor:Magnesium. ,Disease:Defects in RPS6KA3 are the cause of Coffin-Lowry syndrome (CLS) [MIM:303600]; an X-linked dominant disorder characterized by severe mental retardation with facial and digital dysmorphisms , and progressive skeletal deformations. ,enzyme regulation:Activated by multiple phosphorylations on threonine and serine residues. ,Function:Serine/threonine kinase that may play a role in mediating the growth-factor and stress induced activation of the transcription factor CREB. ,PTM:Autophosphorylated on Ser-386 , as part of the activation process. ,PTM:Ser-227 phosphorylation promotes Ser-386 phosphorylation and leads to basal activation. Full activation by growth factors requires additional phosphorylation on Ser-369. ,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily. ,similarity:Contains 1 AGC-kinase C-terminal domain. ,similarity:Contains 2 protein kinase domains. ,subunit:Forms a complex with either ERK1 or ERK2 in quiescent cells. Transiently dissociates following mitogenic stimulation (By similarity) . Interacts with NFATC4. ,tissue specificity:Expressed in many tissues , highest levels in skeletal muscle. ,

| Validation Data



Western Blot analysis of various,using primary antibody at 1:1000 dilution.
Secondary antibody (catalog#:RS23920) was diluted at 1:10000

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**RSK2 (Phospho
Tyr529) Rabbit pAb**