

# Retinoic Acid Receptor $\alpha/\beta$ (Phospho Ser96) Rabbit pAb

CatalogNo: YP1780

## Key Features

### Host Species

- Rabbit

### Reactivity

- Human, Mouse, Rat

### Applications

- WB

### MW

- 50kD (Observed)

### Isotype

- IgG

## Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

## Recommended Dilution Ratios

WB 1:500-2000

## Basic Information

**Clonality** Polyclonal

## Immunogen Information

**Immunogen** Synthesized peptide derived from human Retinoic Acid Receptor  $\alpha/\beta$  (Phospho-Ser96)

**Specificity** This antibody detects endogenous levels of Retinoic Acid Receptor  $\alpha/\beta$  (Phospho-Ser96) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): KSsGY

---

## | Target Information

**Gene name** RARA NR1B1

**Protein Name** Retinoic Acid Receptor  $\alpha/\beta$  (Phospho-Ser96)

Organism	Gene ID	UniProt ID
Human	<a href="#">5914;</a>	<a href="#">P10276;</a>
Mouse	<a href="#">19401;</a>	<a href="#">P11416;</a>

**Cellular Localization** Nucleus . Cytoplasm . Nuclear localization depends on ligand binding, phosphorylation and sumoylation (PubMed:19850744). Translocation to the nucleus in the absence of ligand is dependent on activation of PKC and the downstream MAPK phosphorylation (By similarity). Increased nuclear localization upon pulsatile shear stress (PubMed:28167758). .


**Tissue specificity** Expressed in monocytes.

**Function** Disease:Chromosomal aberrations involving RARA may be a cause of acute promyelocytic leukemia (APL) [MIM:612376]. Translocation t(11;17)(q32;q21) with ZBTB16/PLZF; translocation t(15;17)(q21;q21) with PML; translocation t(5;17)(q32;q11) with NPM.,Domain:Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain.,Function:This is a receptor for retinoic acid. This metabolite has profound effects on vertebrate development. Retinoic acid is a morphogen and is a powerful teratogen. This receptor controls cell function by directly regulating gene expression.,online information:Retinoic acid receptor entry,PTM:Phosphorylated. Phosphorylation does not change during cell cycle. Phosphorylation on Ser-77 is crucial for transcriptional activity.,similarity:Belongs to the nuclear hormone receptor family.,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Interacts with CDK7 (By similarity). Interacts with NCOA3 and NCOA6 coactivators, leading to a strong increase of transcription of target genes. Interacts with NOCA7 in a ligand-inducible manner.,

---

## | Validation Data

Western Blot analysis of various, using primary antibody at 1:1000 dilution.  
Secondary antibody(catalog#:RS23920) was diluted at 1:10000



## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**Retinoic Acid  
Receptor  $\alpha/\beta$   
(Phospho Ser96)  
Rabbit pAb**

