

STYK1/NOK (Phospho Tyr327) Rabbit pAb

CatalogNo: YP1794 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 47kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human STYK1/NOK (Phospho-Tyr327)

Specificity This antibody detects endogenous levels of STYK1/NOK (Phospho-Tyr327) at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): PPyPE

| Target Information

Gene name STYK1 NOK

Protein Name STYK1/NOK (Phospho-Tyr327)


Organism	Gene ID	UniProt ID
Human	55359 ;	Q6J9G0 ;
Mouse	243659 ;	Q6J9G1 ;

Cellular Localization Membrane ; Single-pass membrane protein .

Tissue specificity Widely expressed. Highly expressed in brain , placenta and prostate. Expressed in tumor cells such as hepatoma cells L-02 , cervix carcinoma cells HeLa , ovary cancer cells Ho8910 and chronic myelogenous leukemia cells K-562 , but not in other tumor cells such as epidermoid carcinoma (A-431) . Undetectable in most normal lung tissues , widely expressed in lung cancers.

Function Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Function:Probable tyrosine protein-kinase , which has strong transforming capabilities on a variety of cell lines. When overexpressed , it can also induce tumor cell invasion as well as metastasis in distant organs. May act by activating both MAP kinase and phosphatidylinositol 3'-kinases (PI3K) pathways. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Contains 1 protein kinase domain. ,tissue specificity:Widely expressed. Highly expressed in brain , placenta and prostate. Expressed in tumor cells such as hepatoma cells LO2 , cervix carcinoma cells HeLa , ovary cancer cells Ho8910 and chronic myelogenous leukemia cells K562 , but not in other tumor cells such as epidermoid carcinoma (A431) . Undetectable in most normal lung tissues , widely expressed in lung cancers. ,

| Validation Data



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody (catalog#:RS23920) was diluted at 1:10000

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
STYK1/NOK
(Phospho Tyr327)
Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)