

Met (Phospho Tyr1365) Rabbit pAb

CatalogNo: YP1821

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, WB

MW

- 155kD (Observed)

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human MET (Phospho Tyr1365)

Specificity This antibody detects endogenous levels of MET (Phospho Tyr1365) Rabbit pAb at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): APyPS

| Target Information

Gene name MET

Protein Name Hepatocyte growth factor receptor (HGF receptor) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met)


Organism	Gene ID	UniProt ID
Human	4233 ;	P08581 ;
Mouse		P16056 ;
Rat	24553 ;	P97523 ;

Cellular Localization Membrane; Single-pass type I membrane protein.; [Isoform 3]: Secreted.

Tissue specificity Expressed in normal hepatocytes as well as in epithelial cells lining the stomach , the small and the large intestine. Found also in basal keratinocytes of esophagus and skin. High levels are found in liver , gastrointestinal tract , thyroid and kidney. Also present in the brain. Expressed in metaphyseal bone (at protein level) (PubMed:26637977) .

Function Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Disease:Activation of MET after rearrangement with the TPR gene produces an oncogenic protein. ,Disease:Defects in MET are a cause of hepatocellular carcinoma (HCC) [MIM:114550]. ,Disease:Defects in MET are a cause of hereditary papillary renal carcinoma (HPRC) [MIM:605074]; also known as papillary renal cell carcinoma 2 (RCCP2) . HPRC is a form of inherited kidney cancer characterized by a predisposition to develop multiple , bilateral papillary renal tumors. The pattern of inheritance is consistent with autosomal dominant transmission with reduced penetrance. ,Disease:Defects in MET may be associated with gastric cancer. ,Disease:Genetic variations in MET may be associated with susceptibility to autism type 9 (AUTS9) [MIM:611015]. Autism is a neurodevelopmental disorder characterized by disturbance in language , perception and socialization. The disorder is classically defined by a triad of limited or absent verbal communication , a lack of reciprocal social interaction or responsiveness , and restricted , stereotypical , and ritualized patterns of interests and behavior. ,Domain:The kinase domain is involved in SPSB1 binding. ,Function:Receptor for hepatocyte growth factor and scatter factor. Has a tyrosine-protein kinase activity. Functions in cell proliferation , scattering , morphogenesis and survival. ,online information:C-MET entry ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 Sema domain. ,similarity:Contains 3 IPT/TIG domains. ,subunit:Heterodimer formed of an alpha chain (50 kDa) and a beta chain (145 kDa) which are disulfide linked. Binds PLXNB1 and GRB2. Interacts with SPSB1 , SPSB2 and SPSB4 (By similarity) . Interacts with INPP5D/SHIP1. When phosphorylated at Tyr-1356 , interacts with INPPL1/SHIP2. Interacts with RANBP9 and RANBP10 , as well as SPSB1 , SPSB2 , SPSB3 and SPSB4. SPSB1 binding occurs in the presence and in the absence of HGF , however HGF treatment has a positive effect on this interaction. Interacts with MUC20; prevents interaction with GRB2 and suppresses hepatocyte growth factor-induced cell proliferation. ,

| Validation Data



Western Blot analysis of 1 HeLa cell, HepG2 cell, using primary antibody at 1:1000 dilution. Secondary antibody (catalog#:RS23920) was diluted at 1:10000

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Met (Phospho Tyr1365) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)