

www.immunoway.com.cn

STAT4 (Phospho Ser721) Rabbit pAb

CatalogNo: YP1857 Orthogonal Validated 💽

Key Features

Host Species

Rabbit

ReactivityHuman,Mouse

Applications
• IHC,WB

MW • 86kD (Observed)

Recommended Dilution Ratios

WB 1:500-2000 IHC 1:50-200

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)		
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		

Basic Information

Clonality Polyclonal

Immunogen Information


Immunogen Synthesized peptide derived from human STAT4 (Phospho Ser721)

Specificity This antibody detects endogenous levels of STAT4 (Phospho Ser721) Rabbit pAb at Human, Mouse.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PMsPS

Target Information

Gene name	STAT4			
Protein Name	Signal transducer and activator of transcription 4			
	Organism	Gene ID	UniProt ID	
	Human	<u>6775;</u>	<u>Q14765;</u>	
	Mouse	<u>20849;</u>	<u>P42228;</u>	
Cellular Localization	Cytoplasm. Nucleus. Translocated into the nucleus in response to phosphorylation.			
Function	Disease:Genetic variations in STAT4 are associated with susceptibility to rheumatoid arthritis (RA) [MIM:180300]. Rheumatoid arthritis is a complex, multifactorial disorder. It is one of the most common autoimmune diseases and it is characterized by inflammation of synovial tissue and joint destruction.,Disease:Genetic variations in STAT4 are associated with susceptibility to systemic lupus erythematosus type 11 (SLEB11) [MIM:612253]. Systemic lupus erythematory, and often febrile multisystemic disorder of connective tissue characterized principally by involvement of the skin, joints, kidneys, and serosal membranes. It is thought to represent a failure of the regulatory mechanisms of the autoimmune system.,Function:Carries out a dual Function: signal transduction and activation of transcription. Involved in IL12 signaling.,PTM:Tyrosine phosphorylated. Serine phosphorylation is also required for maximal transcriptional activity.,similarity:Belongs to the transcription factor STAT family.,similarity:Contains 1 SH2 domain.,subcellular location:Translocated into the nucleus in response to phosphorylation.,subunit:Forms a homodimer or a heterodimer with a related family member (By similarity). The SH2 domain interacts, in vitro, with IL12RB2 via a short cytoplasmic domain.			

Validation Data



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

Contact information

- Orders: order.cn@immunoway.com
- Support: support.cn@immunoway.com
- Telephone: 400-8787-807(China)
- Website: http://www.immunoway.com.cn
- Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: STAT4 (Phospho Ser721) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents