

ARA70 Rabbit pAb

CatalogNo: YT0302 Orthogonal Validated 💽

Comparable Abs C

Key Features

Host Species Reactivity Applications

Rabbit
Human, Mouse, Rat
WB, IHC, IF, CoIP, ELISA

MW Isotype • 70kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000

IP 1:50-200Not yet tested in other applications

IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from ARA70 . at AA range: 320-400

Specificity ARA70 Polyclonal Antibody detects endogenous levels of ARA70 protein.

| Target Information

Gene name NCOA4

Protein Name Nuclear receptor coactivator 4

Organism Gene ID UniProt ID

Human <u>8031;</u> <u>Q13772;</u>

Cellular Localization nucleus, autolysosome,

Tissue specificity Widely expressed. Also detected in adipose tissues and in different cell lines. Isoform Beta

is only expressed in testis.

Function Alternative products:Additional isoforms seem to exist,Disease:A chromosomal aberration


involving NCOA4 is a cause of thyroid papillary carcinoma (PACT) [MIM:188550]. Inversion inv(10)(q11.2;q11.2) generates the RET/NCOA4 (PTC3) oncogene that has been found in

sporadic and radiation-associated post-Chernobyl thyroid papillary


carcinomas.,Function:Enhances the androgen receptor transcriptional activity in prostate cancer cells. Ligand-independent coactivator of the peroxisome proliferator-activated receptor (PPAR) gamma.,polymorphism:Polymorphisms are specific to isoform alpha. Not found in isoform beta. Could be encoded by two different genes.,subunit:Interacts with the androgen receptor and the retinoid X receptor (RXR) in a ligand-dependent manner.,tissue specificity:Widely expressed. Also detected in adipose tissues and in different cell lines.

Isoform beta is only expressed in testis.,

Validation Data



Western Blot analysis of various cells using ARA70 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western blot analysis of NCOA4 Antibody. The lane on the right is blocked with the NCOA4 peptide.

| Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **ARA70 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents