

#### www.immunoway.com.cn

# **CREB-1** Rabbit pAb

CatalogNo: YT1097 Orthogonal Validated 💽

## Key Features

Host Species <ul> <li>Rabbit</li> </ul>	Reactivity <ul> <li>Human,Mouse,Rat</li> </ul>	<ul><li>Applications</li><li>WB,IHC,IF,ELISA</li></ul>
MW • 37kD (Observed)	Isotype • IgG	

### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:40000 Not yet tested in other applications.

### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

## Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human CREB. AA range:100-149
Specificity	CREB-1 Polyclonal Antibody detects endogenous levels of CREB-1 protein.

# Target Information

Gene name	CREB1			
Protein Name	Cyclic AMP-responsive element-binding protein 1 Organism Gene ID UniProt ID			
	Human	<u>1385;</u>	<u>P16220;</u>	
	Mouse	<u>12912;</u>	<u>Q01147;</u>	
	Rat	<u>81646;</u>	<u>P15337;</u>	
Cellular Localization	Nucleus .			
Tissue specificity	Eye,Placenta,Spleen,Testis,			
Function	Disease:A chromosomal aberration involving CREB1 is associated with angiomatoid fibrous histiocytoma (AFH) [MIM:612160]. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type.,Function:This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. Transciption activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Implicated in synchronization of circadian rhythmicity.,PTM:Stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the bZIP family.,similarity:Contains 1 bZIP domain.,similarity:Contains 1 KID (kinase-inducible) domain.,subunit:Interacts with PPRC1. Binds DNA as a dimer. This dimer is stabilized by magnesium ions. Interacts, through the bZIP domain, with the coactivators TORC1/CRTC1, TORC2/CRTC2 and TORC3/CRTC3. When phosphorylated on Ser-133, binds CREBBP.,			

# Validation Data

# **Contact information**

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **CREB-1 Rabbit pAb**  For Research Use Only. Not for Use in Diagnostic Procedures.

Immunoway - 3 / 3