

www.immunoway.com.cn

DRP1 Rabbit pAb

CatalogNo: YT1414

Key Features

Host SpeciesReactivity• Rabbit• Human,Mouse,RatMWIsotype• 80kD (Observed)• IgG

Applications • IF,WB,IHC,ELISA

Recommended Dilution Ratios

IF 1:50-200 WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:10000 Not yet tested in other applications

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenSynthesized peptide derived from DRP1 . at AA range: 580-660SpecificityDRP1 Polyclonal Antibody detects endogenous levels of DRP1 protein.

Target Information

Gene name

DNM1L

Protein Name Dynamin-1-like protein

Organism	Gene ID	UniProt ID
Human	<u>10059;</u>	<u>000429;</u>
Mouse	<u>74006;</u>	<u>Q8K1M6;</u>
Rat	<u>114114;</u>	<u>035303;</u>

Cellular Localization

Cytoplasm, cytosol. Golgi apparatus. Endomembrane system; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein. Peroxisome. Membrane, clathrin-coated pit . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane . Mainly cytosolic. Recruited by RALA and RALBP1 to mitochondrion during mitosis (PubMed:21822277). Translocated to the mitochondrial membrane through O-GlcNAcylation and interaction with FIS1. Colocalized with MARCHF5 at mitochondrial membrane. Localizes to mitochondria at sites of division. Localizes to mitochondria following necrosis induction. Recruited to the mitochondrial outer membrane by interaction with MIEF1. Mitochondrial recruitment is inhibited by C11orf65/MFI (By similarity). Associated with peroxisomal membranes, partly recruited there by PEX11B. May also be associated with endoplasmic reticulum tubules and cytoplasmic vesicles and found to be perinuclear. In some cell types, localizes to the Golgi complex (By similarity). Binds to phospholipid membranes (By similarity). .

- **Tissue specificity** Ubiquitously expressed with highest levels found in skeletal muscles, heart, kidney and brain. Isoform 1 is brain-specific. Isoform 2 and isoform 3 are predominantly expressed in testis and skeletal muscles respectively. Isoform 4 is weakly expressed in brain, heart and kidney. Isoform 5 is dominantly expressed in liver, heart and kidney. Isoform 6 is expressed in neurons.
- **Function** Catalytic activity: GTP + H(2)O = GDP + phosphate., Function: Functions in mitochondrial andperoxisomal division probably by regulating membrane fission. Enzyme hydrolyzing GTP that oligomerizes to form ring-like structures and is able to remodel membranes. May also play a role on organelles of the secretory pathway., miscellaneous: Isoform 1 and isoform 2 inhibits peroxisomal division when overexpressed while isoform 3 and isoform 4 have no effect., PTM: Phosphorylated by GSK3B., similarity: Belongs to the dynamin family., similarity: Contains 1 GED domain., subcellular location: Mainly cytosolic. Also membrane-associated. Localizes to mitochondria at spots of division events. Associated with peroxisomal membranes, it is recruited in part by PEX11B. May also be associated with endoplasmic reticulum tubules and cytoplasmic vesicles and found to be perinuclear., subunit: Homotetramer; N-terminal part binds to the C-terminal part of another DNM1L. Can self-assemble in multimeric ring-like structures. Interacts with FIS1 (By similarity). Interacts with GSK3B., tissue specificity: Ubiguitously expressed with highest levels found in skeletal muscles, heart, kidney and brain. Isoform 1 is brain-specific while isoform 3 and isoform 4 are predominantly expressed in testis and skeletal muscles respectively. Isoform 2 is weakly expressed in brain, heart and kidney and isoform 5 is dominantly expressed in liver, heart and kidney.,

Validation Data

Contact information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **DRP1 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents