

## EDG-3 Rabbit pAb

CatalogNo: YT1462

Orthogonal Validated 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IF, ELISA

#### MW

- 42kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-1:2000****IF 1:200-1:1000****ELISA 1:40000****Not yet tested in other applications.**

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

**Clonality** Polyclonal


### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human EDG3. AA range:115-164**Specificity** EDG-3 Polyclonal Antibody detects endogenous levels of EDG-3 protein.


### Target Information

Gene name	S1PR3		
Protein Name	Sphingosine 1-phosphate receptor 3		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">1903</a> ;	<a href="#">Q99500</a> ;
	Mouse	<a href="#">13610</a> ;	<a href="#">Q9Z0U9</a> ;
Cellular Localization	Cell membrane; Multi-pass membrane protein.		
Tissue specificity	Expressed in all tissues, but most abundantly in heart, placenta, kidney, and liver.		
Function	Function:Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. When expressed in rat HTC4 hepatoma cells, is capable of mediating S1P-induced cell proliferation and suppression of apoptosis.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in all tissues, but most abundantly in heart, placenta, kidney, and liver.,		


Validation Data



Yuan Chang, Hu, et al. "Hyperglycemia Triggered S1P/S1PR3 Signaling Worsens Liver Ischemia/Reperfusion Injury by Regulating M1/M2 Polarization." Chao and Yang, Shikun and Cheng, Xuyu and Cheng, Feng and Rao, Jianhua and Wang, Xue-Hao, Hyperglycemia Triggered S1P/S1PR3 Signaling Worsens Liver Ischemia/Reperfusion Injury by Regulating M 1 (2018).



Western Blot analysis of HT-29 cells using EDG-3 Polyclonal Antibody



Western blot analysis of lysates from HT-29 cells, using EDG3 Antibody. The lane on the right is blocked with the synthesized peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
 Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**EDG-3 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)