**Applications** 

WB,ELISA



# **Elf-4 Rabbit pAb**

CatalogNo: YT1524

# **Key Features**

Host Species Reactivity

RabbitHuman, Mouse

MW Isotype

• 70kD (Observed) • IgG

### **Recommended Dilution Ratios**

WB 1:500-1:2000 ELISA 1:10000

Not yet tested in other applications.

### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

**Clonality** Polyclonal

## Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human ELF4. AA

range:178-227

**Specificity** Elf-4 Polyclonal Antibody detects endogenous levels of Elf-4 protein.

# | Target Information

#### Gene name

ELF4

#### **Protein Name**

ETS-related transcription factor Elf-4

Organism	Gene ID	UniProt ID
Human	<u>2000</u> ;	<u>Q99607;</u>
Mouse	<u>56501</u> ;	<u>Q9Z2U4;</u>

### Cellular Localization


Nucleus, PML body. Accumulation into PML nuclear bodies is mediated by PML.

**Tissue specificity** Abundantly expressed in the placenta and in a variety of myeloid leukemia cell lines. Moderate levels of expression in heart, lung, spleen, thymus, peripheral blood lymphocytes, ovary and colon. Lower levels of expression in Jurkat T-cells and other T-cell lines and no expression in brain.


#### **Function**

Function: Transcriptional activator that binds to DNA sequences containing the consensus 5'-WGGA-3'. Transactivates promoters of the hematopoietic growth factor genes CSF2, IL3, IL8, and of the bovine lysozyme gene. Acts synergistically with RUNX1 to transactivate the IL3 promoter. Also transactivates the PRF1 promoter in natural killer (NK) cells. Plays a role in the development and function of NK and NK T-cells and in innate immunity..induction:By ponisterone A in erythroleukemia cells., similarity: Belongs to the ETS family, similarity: Contains 1 ETS DNA-binding domain, subcellular location: Accumulation into PML nuclear bodies is mediated by PML, subunit: Interacts via an N-terminal region with the RUNX1 Runt domain. Interacts via its C-terminus with PML; this interaction enhances ELF4 transactivation of the bovine lysozyme gene., tissue specificity: Abundantly expressed in the placenta and in a variety of myeloid leukemia cell lines. Moderate levels of expression in heart, lung, spleen, thymus, peripheral blood lymphocytes, ovary and colon. Lower levels of expression in Jurkat T-cells and other T-cell lines and no expression in brain.,

# **I** Validation Data



Western Blot analysis of 22RV1 cells using Elf-4 Polyclonal Antibody diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western blot analysis of the lysates from RAW264.7cells using ELF4 antibody.

# | Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Elf-4 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents