

FoxO1A Rabbit pAb

CatalogNo: YT1761

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 70kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:10000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human FOXO1A. AA range:295-344

Specificity FoxO1A Polyclonal Antibody detects endogenous levels of FoxO1A protein.

| Target Information

Gene name FOXO1

Protein Name Forkhead box protein O1

Organism	Gene ID	UniProt ID
Human	2308 ;	Q12778 ;
Mouse	56458 ;	Q9R1E0 ;
Rat	84482 ;	G3V7R4 ;

Cellular Localization Cytoplasm . Nucleus . Shuttles between the cytoplasm and nucleus. Largely nuclear in unstimulated cells (PubMed:11311120 , PubMed:12228231 , PubMed:19221179 , PubMed:21245099 , PubMed:20543840 , PubMed:25009184) . In osteoblasts , colocalizes with ATF4 and RUNX2 in the nucleus (By similarity) . Serum deprivation increases localization to the nucleus , leading to activate expression of SOX9 and subsequent chondrogenesis (By similarity) . Insulin-induced phosphorylation at Ser-256 by PKB/AKT1 leads , via stimulation of Thr-24 phosphorylation , to binding of 14-3-3 proteins and nuclear export to the cytoplasm where it is degraded by the ubiquitin-proteosomal pathway (PubMed:11237865 , PubMed:12228231) . Phosphorylation at Ser-249 by CDK1 disrupts binding of 14-3-3 proteins and promotes nuclear accumulation (PubMed:18356527) . Phosphorylation by NLK results in nuclear export (By similarity) . Translocates to the nucleus upon oxidative stress-induced phosphorylation at Ser-212 by STK4/MST1 (PubMed:19221179 , PubMed:21245099) . SGK1-mediated phosphorylation also results in nuclear translocation (By similarity) . Retained in the nucleus under stress stimuli including oxidative stress , nutrient deprivation or nitric oxide (By similarity) . Retained in the nucleus on methylation (By similarity) . PPIA/CYPA stimulates its nuclear accumulation (PubMed:31063815) . Deacetylation by SIRT6 , promotes its translocation into the cytoplasm (PubMed:25009184) . .

Tissue specificity Ubiquitous.

Function Disease:Chromosomal aberrations involving FOXO1 are a cause of rhabdomyosarcoma 2 (RMS2) [MIM:268220]; also known as alveolar rhabdomyosarcoma. Translocation (2;13) (q35;q14) with PAX3; translocation t (1;13) (p36;q14) with PAX7. The resulting protein is a transcriptional activator. ,Function:Transcription factor. ,PTM:Phosphorylated by AKT1; insulin-induced (By similarity) . IGF1 rapidly induces phosphorylation of Ser-256 , Thr-24 , and Ser-319. Phosphorylation of Ser-256 decreases DNA-binding activity and promotes the phosphorylation of Thr-24 , and Ser-319 , permitting phosphorylation of Ser-322 and Ser-325 , probably by CK1 , leading to nuclear exclusion and loss of function. Phosphorylation of Ser-329 is independent of IGF1 and leads to reduced function. Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity:Contains 1 fork-head DNA-binding domain. ,subcellular location:Shuttles between cytoplasm and nucleus. ,subunit:Interacts with LRPPRC. ,tissue specificity:Ubiquitous. ,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
FoxO1A Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)