

# Gab 2 Rabbit pAb

CatalogNo: YT1815 Orthogonal Validated (9)

## **Key Features**

**Host Species** 

Reactivity Rabbit

Human, Mouse, Rat

**Applications** 

WB,IHC,IF,ELISA

MW 74kD (Observed) Isotype IgG

## Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 **ELISA 1:40000** IF 1:50-200

## Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

## **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

The antiserum was produced against synthesized peptide derived from human Gab2. AA **Immunogen** 

range:589-638

**Specificity** Gab 2 Polyclonal Antibody detects endogenous levels of Gab 2 protein.

# | Target Information

#### Gene name

GAB2

### **Protein Name**

GRB2-associated-binding protein 2

| Organism | Gene ID       | UniProt ID     |
|----------|---------------|----------------|
| Human    | <u>9846;</u>  | Q9UQC2;        |
| Mouse    |               | <u>Q9Z1S8;</u> |
| Rat      | <u>84477;</u> | Q9EQH1;        |

### Cellular Localization


Cytoplasm . Cell membrane .

Tissue specificity Brain, Clones donated by Kazusa DNA Research Inst.,


### **Function**

PTM:Dephosphorylated by PTPN11.,PTM:Phosphorylated on tyrosine residue(s) by the thrombopoietin receptor (TPOR), stem cell factor receptor (SCFR), and T-cell and B-cell antigen receptors, gp130, IL-2R and IL-3R., similarity: Belongs to the GAB family., similarity: Contains 1 PH domain., subunit: Interacts with GRB2, PI-3 kinase and with other SH2-containing proteins.,

### **Validation Data**



Western blot analysis of lysates from Jurkat cells, treated with TNF 2500U/ml 30', using Gab2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from RAW264.7cells using Gab2 antibody.



Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200

### **|** Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Gab 2 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents