

HBO1 Rabbit pAb

CatalogNo: Y2101

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA

MW

- 75kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human MYST2. AA range:131-180

Specificity HBO1 Polyclonal Antibody detects endogenous levels of HBO1 protein.

Target Information

Gene name KAT7

Protein Name Histone acetyltransferase KAT7

Organism	Gene ID	UniProt ID
Human	11143 ;	O95251 ;
Mouse	217127 ;	Q5SVQ0 ;
Rat	303470 ;	Q810T5 ;

Cellular Localization

Nucleus . Chromosome . Chromosome , centromere . Cytoplasm , cytosol . Associates with replication origins specifically during the G1 phase of the cell cycle (PubMed:18832067 , PubMed:20129055) . Localizes to transcription start sites (PubMed:21753189 , PubMed:24065767) . Localizes to ultraviolet-induced DNA damage sites following phosphorylation by ATR (PubMed:28719581) . Localizes to centromeres in G1 phase (PubMed:27270040) . .

Tissue specificity Ubiquitously expressed , with highest levels in testis.

Function

Catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone. ,Domain:The C2HC-type zinc finger is required for interaction with MCM2 and ORC1L. ,Domain:The N-terminus is involved in transcriptional repression , while the C-terminus mediates AR-interaction. ,Function:Component of the HBO1 complex which has a histone H4-specific acetyltransferase activity , a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription. ,PTM:Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity:Belongs to the MYST (SAS/MOZ) family. ,similarity:Contains 1 C2HC-type zinc finger. ,subunit:Component of the HBO1 complex composed at least of ING4 or ING5 , MYTS2/HBO1 , EAF6 , and one of PHF15 , PHF16 and PHF17. Interacts with MCM2 and ORC1L. Interacts with the androgen receptor (AR) in the presence of dihydrotestosterone. ,tissue specificity:Ubiquitously expressed , with highest levels in testis. ,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
HBO1 Rabbit pAb