

## hnRNP K Rabbit pAb

CatalogNo: YT2202

### **Key Features**

**Host Species** 

Rabbit

Reactivity

· Human, Mouse, Rat

Applications
• WB,IHC,IF,ELISA

MW

51kD (Observed)

IsotypeIgG

### Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000 IF 1:50-200

### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human hnRNP K.

AA range:250-299

**Specificity** hnRNP K Polyclonal Antibody detects endogenous levels of hnRNP K protein.

# | Target Information

#### Gene name HNRNPK

#### **Protein Name**

Heterogeneous nuclear ribonucleoprotein K

Organism	Gene ID	UniProt ID
Human	<u>3190;</u>	<u>P61978;</u>
Mouse	<u>15387;</u>	<u>P61979;</u>
Rat	<u>117282;</u>	<u>P61980</u> ;

#### Cellular Localization


Cytoplasm . Nucleus, nucleoplasm . Cell projection, podosome . Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear (PubMed:18775702).

**Tissue specificity** Brain, Cajal-Retzius cell, Colorectal cancer and surrounding


#### **Function**

Function:One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for premRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA., mass spectrometry: PubMed:11840567, PTM: Arg-296 and Arg-299 are dimethylated, probably to asymmetric dimethylarginine., similarity: Contains 1 KH domain., similarity: Contains 2 KH domains., similarity: Contains 3 KH domains., subcellular location:In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear., subunit: Interacts with RBM42 and ZIK1 (By similarity). Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRNPK, HNRPM, HNRPR, HNRPU, KIAA1160, KIAA1604, LSM2, LSM3, MAGOH, MORG1, PABPC1, PLRG1, PNN, PPIE. PPIL1, PPIL3, PPWD1, PRPF19, PRPF4B, PRPF6, PRPF8, RALY, RBM22, RBM8A, RBMX, SART1, SF3A1, SF3A2, SF3A3, SF3B1, SF3B2, SF3B3, SFRS1, SKIV2L2, SNRPA1, SNRPB, SNRPB2, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF, SNRPG, SNW1, SRRM1, SRRM2, SYF2, SYNCRIP, TFIP11, THOC4, U2AF1, WDR57, XAB2 and ZCCHC8. Interacts with ANKRD28. Interacts with HCV core protein. Interacts with ASFV p30 protein.,


### **Validation Data**



Western Blot analysis of various cells using hnRNP K Polyclonal Antibody diluted at 1:2000



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from K562, A549, and HT-29 cells, using hnRNP K Antibody. The lane on the right is blocked with the synthesized peptide.

### **Contact information**

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

hnRNP K Rabbit pAb