

MaxiK β Rabbit pAb

CatalogNo: YT2667

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 24kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

The antiserum was produced against synthesized peptide derived from human MaxiK β . AA range: 70-119


Specificity

MaxiK β Polyclonal Antibody detects endogenous levels of MaxiK β protein.


Target Information

Gene name	KCNMB4		
Protein Name	Calcium-activated potassium channel subunit beta-4		
	Organism	Gene ID	UniProt ID
	Human	27345;	Q86W47;
	Mouse	58802;	Q9JIN6;
	Rat	66016;	Q9ESK8;
Cellular Localization	Membrane; Multi-pass membrane protein.		
Tissue specificity	Predominantly expressed in brain. In brain, it is expressed in the cerebellum, cerebral cortex, medulla, spinal cord, occipital pole, frontal lobe, temporal lobe, putamen, amygdala, caudate nucleus, corpus callosum, hippocampus, substantia nigra and thalamus. Weakly or not expressed in other tissues.		
Function	<p>Domain:Resistance to charybdotoxin (CTX) toxin is mediated by the extracellular domain.,Function:Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel. Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to KCNMA1 channel diversity. Decreases the gating kinetics and calcium sensitivity of the KCNMA1 channel, but with fast deactivation kinetics. May decrease KCNMA1 channel openings at low calcium concentrations but increases channel openings at high calcium concentrations. Makes KCNMA1 channel resistant to 100 nM charybdotoxin (CTX) toxin concentrations.,miscellaneous:Treatment with okadaic acid reduces its effect on KCNMA1.,PTM:N-glycosylated. A highly glycosylated form is promoted by KCNMA1. Glycosylation, which is not required for the interaction with KCNMA1 and subcellular location, increases protection against charybdotoxin.,PTM:Phosphorylated. Phosphorylation modulates its effect on KCNMA1 activation kinetics.,similarity:Belongs to the KCNMB family.,subunit:Interacts with KCNMA1 tetramer. There are probably 4 molecules of KCMNB4 per KCNMA1 tetramer.,tissue specificity:Predominantly expressed in brain. In brain, it is expressed in the cerebellum, cerebral cortex, medulla, spinal cord, occipital pole, frontal lobe, temporal lobe, putamen, amygdala, caudate nucleus, corpus callosum, hippocampus, substantia nigra and thalamus. Weakly or not expressed in other tissues.,</p>		


| Validation Data



Western Blot analysis of various cells using MaxiK β Polyclonal Antibody



Immunohistochemistry analysis of MaxiK β antibody in paraffin-embedded human brain tissue.



Western blot analysis of lysate from HepG2 cells, using MaxiK β antibody.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
 to access additional
 product information:
MaxiK β Rabbit pAb

