

## Mena Rabbit pAb

CatalogNo: YT2731

### | Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 67kD (Observed)

#### Isotype

- IgG

### | Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:10000**

**IF 1:50-200**

### | Storage

#### Storage\*

-15°C to -25°C/1 year (Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### | Basic Information

#### Clonality

Polyclonal

### | Immunogen Information

#### Immunogen

The antiserum was produced against synthesized peptide derived from human ENAH. AA range: 472-521


#### Specificity

Mena Polyclonal Antibody detects endogenous levels of Mena protein.


### | Target Information

Gene name	ENAH		
Protein Name	Protein enabled homolog		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">55740;</a>	<a href="#">Q8N8S7;</a>
	Mouse	<a href="#">13800;</a>	<a href="#">Q03173;</a>
Cellular Localization	Cytoplasm. Cytoplasm, cytoskeleton . Cell projection, lamellipodium . Cell projection, filopodium . Cell junction, synapse . Cell junction, focal adhesion. Targeted to the leading edge of lamellipodia and filopodia by MRL family members. Colocalizes at filopodial tips with a number of other proteins including vinculin and zyxlin. Colocalizes with N-WASP at the leading edge. Colocalizes with GPHN and PFN at synapses (By similarity). .		
Tissue specificity	Expressed in myoepithelia of parotid, breast, bronchial glands and sweat glands. Expressed in colon-rectum muscircularis mucosae epithelium, pancreas acinar ductal epithelium, endometrium epithelium, prostate fibromuscular stroma and placenta vascular media. Overexpressed in a majority of breast cancer cell lines and primary breast tumor lesions.		
Function	<p>Domain:The EVH2 domain is comprised of 3 regions. Block A is a thymosin-like domain required for G-actin binding. The KLKR motif within this block is essential for the G-actin binding and for actin polymerization. Block B is required for F-actin binding and subcellular location, and Block C for tetramerization.,Function:Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance and lamellipodial and filopodial dynamics in migrating cells. ENAH induces the formation of F-actin rich outgrowths in fibroblasts. Acts syngeristically with BAIAP2-alpha and downstream of NTN1 to promote filopodia formation. Required for the actin-based mobility of Listeria monocytogenes.,PTM:NTN1-induced PKA phosphorylation on Ser-265 directly parallels the formation of filopodial protrusions.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the Ena/VASP family.,similarity:Contains 1 WH1 domain.,subcellular location:Targeted to the leading edge of lamellipodia and filopodia by MRL family members. Colocalizes at filopodial tips with a number of other proteins including vinculin and zyxlin. Colocalizes with N-WASP at the leading edge. Colocalizes with GPHN and PFN at synapses.,subunit:Homotetramer (By similarity). Interacts with APBB1IP, PFN1 and ROBO4. Isoforms, containing the polyproline-rich regions with PPLP motifs, bind the WW domain of APBB1IP. Isoforms, containing the PPSY motif, bind, in vitro, to the WW2 and WW3 domains of NEDD4 and to the WW1 domain of YAP1. Binds the SH3 domain of BAIAP2-alpha but only after the autoinhibitory region of BAIAP2-alpha has been blocked by interaction with CDC42. Interacts, via the EVH1/WH1 domain, with the Pro-rich domains from VCL, ZYX and Listeria monocytogenes actA. Interaction with ZYX is important for targeting ENAH to focal adhesions and enhances production of actin-rich structures at the apical surface of cells. Interacts, through the Pro-rich region, with the C-terminal SH3 domain of DNMPB. Binds GPHN.,tissue specificity:Expressed in myoepithelia of parotid, breast, bronchial glands and sweat glands. Expressed in colon-rectum muscircularis mucosae epithelium, pancreas acinar ductal epithelium, endometrium epithelium, prostate fibromuscular stroma and placenta vascular media. Overexpressed in a majority of breast cancer cell lines and primary breast tumor lesions.,</p>		


| Validation Data



Western Blot analysis of 293 cells using Mena Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using ENAH Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 cells, using ENAH Antibody. The lane on the right is blocked with the synthesized peptide.

## Contact information

Orders: order.cn@immunoway.com  
 Support: support.cn@immunoway.com  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**Mena Rabbit pAb**