

## mTOR Rabbit pAb

CatalogNo: YT2913 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat, Bovine, Chicken, Pig

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 289kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-2000**

**IHC 1:100-500**

**IF/ICC 1:100-500**

**ELISA 1:5000-20000**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human mTOR. AA range:2447-2496

**Specificity** mTOR Polyclonal Antibody detects endogenous levels of mTOR protein.

## Target Information

**Gene name** MTOR

**Protein Name** Serine/threonine-protein kinase mTOR

Organism	Gene ID	UniProt ID
Human	<a href="#">2475;</a>	<a href="#">P42345;</a>
Mouse	<a href="#">56717;</a>	<a href="#">Q9JLN9;</a>
Rat	<a href="#">56718;</a>	<a href="#">P42346;</a>

### Cellular Localization

Endoplasmic reticulum membrane ; Peripheral membrane protein ; Cytoplasmic side . Golgi apparatus membrane ; Peripheral membrane protein ; Cytoplasmic side . Mitochondrion outer membrane ; Peripheral membrane protein ; Cytoplasmic side . Lysosome . Cytoplasm . Nucleus , PML body . Microsome membrane . Lysosome membrane . Cytoplasmic vesicle , phagosome . Shuttles between cytoplasm and nucleus. Accumulates in the nucleus in response to hypoxia (By similarity) . Targeting to lysosomes depends on amino acid availability and RRAGA and RRAGB (PubMed:18497260 , PubMed:20381137) . Lysosome targeting also depends on interaction with MEAK7. Translocates to the lysosome membrane in the presence of TM4SF5 (PubMed:30956113) . .

**Tissue specificity** Expressed in numerous tissues , with highest levels in testis.

### Function

Function:Acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. Part of the TORC2 complex which plays a critical role in AKT1 Ser-473 phosphorylation , and may modulate the phosphorylation of PKCA and regulate actin cytoskeleton organization. ,similarity:Belongs to the PI3/PI4-kinase family. ,similarity:Contains 1 FAT domain. ,similarity:Contains 1 FATC domain. ,similarity:Contains 1 PI3K/PI4K domain. ,similarity:Contains 7 HEAT repeats. ,subunit:Interacts with the FKBP12-rapamycin complex. Binds UBQLN1. Forms part of the mammalian target of rapamycin 2 complex (TORC2) comprised of FRAP1 , GBL , PRR5 , RICTOR and SIN. TORC2 does not bind to and is not sensitive to FKBP12-rapamycin. Binds directly to PRR5 and RICTOR within the TORC2 complex. ,tissue specificity:Expressed in numerous tissues , with highest levels in testis. ,

## Validation Data

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**mTOR Rabbit pAb**

