ApplicationsWB,IHC,IF,ELISA



NGF Rabbit pAb

CatalogNo: YT3114 Orthogonal Validated 💽

Key Features

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype • 27kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-1:2000 IHC: 1:100-300 ELISA 1:20000 IF 1:100-300

Not yet tested in other applications.

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human NGF. AA

range:33-82

Specificity NGF Polyclonal Antibody detects endogenous levels of NGF protein.

| Target Information

Gene name

NGF

Protein Name

Beta-nerve growth factor

Organism	Gene ID	UniProt ID
Human	<u>4803;</u>	<u>P01138;</u>
Mouse	<u>18049</u> ;	<u>P01139</u> ;
Rat	<u>310738;</u>	<u>P25427;</u>


Cellular Localization Secreted . Endosome lumen . ProNGF is endocytosed after binding to the cell surface receptor formed by SORT1 and NGFR. .

Tissue specificity Brain, Epithelium, Eye, Leukocyte,


Function

Disease: Defects in NGF are the cause of hereditary sensory and autonomic neuropathy type 5 (HSAN5) [MIM:608654]. The hereditary sensory and autonomic neuropathies are a genetically and clinically heterogeneous group of disorders characterized by degeneration of dorsal root and autonomic ganglion cells, and by sensory and/or autonomic abnormalities. HSAN5 patients manifest loss of pain perception and impaired temperature sensitivity, ulcers, and in some cases self-mutilation. The autonomic involvement is variable. Function: Nerve growth factor is important for the development and maintenance of the sympathetic and sensory nervous systems. It stimulates division and differentiation of sympathetic and embryonic sensory neurons., online information: Nerve growth factor entry, similarity: Belongs to the NGF-beta family., subunit: Homodimer.,


Validation Data




Immunofluorescence analysis of A549. 1, primary Antibody (red) was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, DAPI(blue) 10min.



Western Blot analysis of Jurkat cells using NGF Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of NGF antibody in paraffin-embedded human lung carcinoma tissue.



Western blot analysis of lysate from Jurkat cells, using NGF antibody.

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **NGF Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody ELISA Kits Protein Reagents
	Immunoway - 4 / 4