

## Peroxin 7 Rabbit pAb

CatalogNo: YT3678

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 40kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:40000**

**IF 1:50-200**

### Storage

#### Storage\*

-15°C to -25°C/1 year (Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

#### Clonality

Polyclonal

### Immunogen Information

#### Immunogen

The antiserum was produced against synthesized peptide derived from human PEX7. AA range: 204-253


#### Specificity

Peroxin 7 Polyclonal Antibody detects endogenous levels of Peroxin 7 protein.


### Target Information

Gene name	PEX7		
Protein Name	Peroxisomal targeting signal 2 receptor		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">5191</a> ;	<a href="#">O00628</a> ;
	Mouse	<a href="#">18634</a> ;	<a href="#">P97865</a> ;
Cellular Localization	Peroxisome . Cytoplasm .		
Tissue specificity	Ubiquitous. Highest expression in pancreas, skeletal muscle and heart.		
Function	<p>Disease:Defects in PEX7 are a cause of Refsum disease (RD) [MIM:266500]; also known as phytanic acid oxidase deficiency. RD is clinically characterized by a tetrad of abnormalities: retinitis pigmentosa, peripheral neuropathy, cerebellar ataxia, and elevated protein levels in the cerebrospinal fluid (CSF). Patients exhibit accumulation of the branched-chain fatty acid, phytanic acid, in blood and tissues. Less constant features are nerve deafness, anosmia, skeletal abnormalities, ichthyosis, cataracts and cardiac impairment. Manifestations of the disease appear in the second or third decade of life.,Disease:Defects in PEX7 are the cause of peroxisome biogenesis disorder complementation group 11 (PBD-CG11) [MIM:601757]. PBD refers to a group of peroxisomal disorders arising from a failure of protein import into the peroxisomal membrane or matrix. The PBD group is comprised of four disorders: Zellweger syndrome (ZWS), neonatal adrenoleukodystrophy (NALD), infantile Refsum disease (IRD), and classical rhizomelic chondrodysplasia punctata (RCDP). ZWS, NALD and IRD are distinct from RCDP and constitute a clinical continuum of overlapping phenotypes known as the Zellweger spectrum. The PBD group is genetically heterogeneous with at least 13 distinct genetic groups as concluded from complementation studies.,Disease:Defects in PEX7 are the cause of rhizomelic chondrodysplasia punctata type 1 (RCDP1) [MIM:215100]. RCDP1 is characterized by rhizomelic shortening of femur and humerus, vertebral disorders, cataract, cutaneous lesions and severe mental retardation.,Function:Binds to the N-terminal PTS2-type peroxisomal targeting signal and plays an essential role in peroxisomal protein import.,similarity:Belongs to the WD repeat peroxin-7 family.,similarity:Contains 6 WD repeats.,subunit:Interacts with PEX5.,tissue specificity:Ubiquitous. Highest expression in pancreas, skeletal muscle and heart.,</p>		


| Validation Data



Western Blot analysis of 3T3 cells using Peroxin 7 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PEX7 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, using PEX7 Antibody. The lane on the right is blocked with the synthesized peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
 Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**Peroxin 7 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)