

## ROCK1 Rabbit pAb

CatalogNo: YT4162

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat, Monkey

#### Applications

- IF, WB, IHC, ELISA

#### MW

- 158kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**IF 1:50-200**

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ICC 1:200-1:1000**

**ELISA 1:20000**

**Not yet tested in other applications**

### Storage

#### Storage\*

-15°C to -25°C/1 year(Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

#### Clonality

Polyclonal

### Immunogen Information

#### Immunogen

The antiserum was produced against synthesized peptide derived from human Rock-1. AA range:262-311

#### Specificity

Rock-1 Polyclonal Antibody detects endogenous levels of Rock-1 protein.

## | Target Information

**Gene name** ROCK1

**Protein Name** Rho-associated protein kinase 1

Organism	Gene ID	UniProt ID
Human	<a href="#">6093</a> ;	<a href="#">Q13464</a> ;
Mouse	<a href="#">19877</a> ;	<a href="#">P70335</a> ;
Rat	<a href="#">81762</a> ;	<a href="#">Q63644</a> ;

**Cellular Localization** Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole . Golgi apparatus membrane ; Peripheral membrane protein . Cell projection, bleb . Cytoplasm, cytoskeleton . Cell membrane . Cell projection, lamellipodium . Cell projection, ruffle . A small proportion is associated with Golgi membranes (PubMed:12773565). Associated with the mother centriole and an intercentriolar linker (By similarity). Colocalizes with ITGB1BP1 and ITGB1 at the cell membrane predominantly in lamellipodia and membrane ruffles, but also in retraction fibers (By similarity). Localizes at the cell membrane in an ITGB1BP1-dependent manner (By similarity). .

**Tissue specificity** Detected in blood platelets.

**Function** Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,Domain:The C-terminal auto-inhibitory domain interferes with kinase activity. RHOA binding leads to a conformation change and activation of the kinase. Truncated ROCK1 is constitutively activated.,enzyme regulation:Activated by RHOA binding.,Function:Protein kinase that phosphorylates a large number of important signaling proteins, and thereby regulates the assembly of the actin cytoskeleton, cell migration, invasiveness of tumor cells, smooth muscle contraction and neurite outgrowth. Necessary for apoptotic membrane blebbing. Plays a role in smooth muscle contraction. Required for centromere positioning and centromere-dependent exit from mitosis.,miscellaneous:Inhibited by Y-27632.,PTM:Autophosphorylated on serine and threonine residues. Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Cleaved by caspase-3 during apoptosis. This leads to constitutive activation of the kinase and membrane blebbing.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 phorbol-ester/DAG-type zinc finger.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 REM (Hr1) repeat.,subcellular location:Associated with the mother centriole and an intercentriolar linker (By similarity). A small proportion is associated with Golgi membranes.,subunit:Binds RHOA (activated by GTP). Interacts with ADD1, GEM, RHOB, RHOC, MYLC2B and VIM (By similarity). Binds RHOE, PPP1R12A, LIMK1 and LIMK2. Interacts with TSG101.,tissue specificity:Detected in blood platelets.,

## | Validation Data

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**ROCK1 Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)