

Siah-2 Rabbit pAb

CatalogNo: YT4297

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 38kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-1:2000**IHC 1:100-1:300****ELISA 1:20000****IF 1:50-200**

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

The antiserum was produced against synthesized peptide derived from human SIAH2. AA range: 241-290


Specificity

Siah-2 Polyclonal Antibody detects endogenous levels of Siah-2 protein.


Target Information

Gene name	SIAH2		
Protein Name	E3 ubiquitin-protein ligase SIAH2		
	Organism	Gene ID	UniProt ID
	Human	6478;	O43255;
	Mouse	20439;	Q06986;
	Rat	140593;	Q8R4T2;
Cellular Localization	Cytoplasm . Nucleus . Predominantly cytoplasmic. Partially nuclear. .		
Tissue specificity	Widely expressed at low level.		
Function	<p>Domain:The RING-type zinc finger domain is essential for ubiquitin ligase activity.,Domain:The SBD domain (substrate-binding domain) mediates the homodimerization and the interaction with substrate proteins. It is related to the TRAF family.,Function:E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Mediates E3 ubiquitin ligase activity either through direct binding to substrates or by functioning as the essential RING domain subunit of larger E3 complexes. Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (POU2AF1, PML, NCOR1), a cell surface receptor (DCC), an antiapoptotic protein (BAG1), and a protein involved in synaptic vesicle function in neurons (SYP). It is thereby involved in apoptosis, tumor suppression, cell cycle, transcription and signaling processes. Has some overlapping function with SIAH1. Triggers the ubiquitin-mediated degradation of TRAF2, whereas SIAH1 can not.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the SINA (Seven in absentia) family.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 SIAH-type zinc finger.,subcellular location:Predominantly cytoplasmic (Probable). Partially nuclear.,subunit:Homodimer. Interacts with UBE2E2. Interacts with PEG3 (By similarity). Interacts with VAV1, without mediating its ubiquitin-mediated degradation. Interacts with CACYBP/SIP. Probable component of some large E3 complex possibly composed of UBE2D1, SIAH2, CACYBP/SIP, SKP1A, APC and TBL1X. Interacts with PEG10, which may inhibit its activity.,tissue specificity:Widely expressed at low level.,</p>		


| Validation Data



Western Blot analysis of various cells using Siah-2 Polyclonal Antibody



Western blot analysis of lysates from COLO cells, using SIAH2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Siah-2 Rabbit pAb

