

TDE1 Rabbit pAb

CatalogNo: YT4591

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 52kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human SERC3. AA range: 361-410

Specificity TDE1 Polyclonal Antibody detects endogenous levels of TDE1 protein.

Target Information

Gene name SERINC3

Protein Name Serine incorporator 3

Organism	Gene ID	UniProt ID
Human	10955 ;	Q13530 ;
Mouse	26943 ;	Q9QZ19 ;


Cellular Localization

Cell membrane ; Multi-pass membrane protein . Golgi apparatus membrane ; Multi-pass membrane protein . ; Cytoplasm, perinuclear region . (Microbial infection) Upon HIV-1 infection, it is redirected to perinuclear region following interaction with HIV-1 Nef, excluding it from virions particles, thereby preventing subsequent antiviral defense activity (Probable). .


Tissue specificity Ubiquitous. Expression levels were increased fourfold to tenfold in lung tumor tissues compared with normal pulmonary tissues.

Function Function:May be involved in cellular transformation.,PTM:N-glycosylated.,similarity:Belongs to the TDE1 family.,tissue specificity:Ubiquitous. Expression levels were increased fourfold to tenfold in lung tumor tissues compared with normal pulmonary tissues.,


Validation Data




Western Blot analysis of various cells using TDE1 Polyclonal Antibody



Immunofluorescence analysis of HepG2 cells, using SERC3 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using SERC3 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
TDE1 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)