

## Nectin 3 Rabbit pAb

CatalogNo: YT5261

### Key Features

60kD (Observed)

Host Species

Rabbit

MW

ReactivityHuman,Mouse,RatIsotype

• IgG

ApplicationsWB,ELISA

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 ELISA 1:20000 Not yet tested in other applications.

#### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human PVRL3. AA range:311-360
Specificity	Nectin 3 Polyclonal Antibody detects endogenous levels of Nectin 3 protein.

#### Target Information

Gene name	VRL3
-----------	------

#### **Protein Name** Poliovirus receptor-related protein 3


Organism	Gene ID	UniProt ID
Human	<u>25945;</u>	<u>Q9NQS3;</u>
Mouse	<u>58998;</u>	<u>Q9JLB9;</u>

CellularCell membrane ; Single-pass membrane protein . Cell junction, synapse, postsynaptic cellLocalizationmembrane .


# **Tissue specificity** Predominantly expressed in testis and placenta as well as in many cell lines, including epithelial cell lines.

FunctionFunction:Plays a role in cell-cell adhesion through heterophilic trans-interactions with<br/>nectin-like proteins or nectins, such as trans-interaction with PVRL2/nectin-2 at Sertoli-<br/>spermatid junctions. Trans-interaction with PVR induces activation of CDC42 and RAC small<br/>G proteins through common signaling molecules such as SRC and RAP1. Also involved in the<br/>formation of cell-cell junctions, including adherens junctions and synapses. Induces<br/>endocytosis-mediated down-regulation of PVR from the cell surface, resulting in reduction of<br/>cell movement and proliferation. Plays a role in the morphology of the ciliary<br/>body.,similarity:Belongs to the nectin family.,similarity:Contains 1 Ig-like V-type<br/>(immunoglobulin-like) domain.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like)<br/>domains.,subunit:Cis- and trans-homodimer. Can form trans-heterodimers with<br/>PVRL1/nectin-1, PVRL2/nectin-2, PVR, IGSF4B/Necl-1 and with IGSF4. Interacts with<br/>MLLT4/afadin. Binds with low affinity to TIGIT.,tissue specificity:Predominantly expressed in<br/>testis and placenta as well as in many cell lines, including epithelial cell lines.,

#### Validation Data



Western Blot analysis of HeLa cells using Nectin 3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from HeLa cells, using PVRL3 Antibody.

#### **Contact information**

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Nectin 3 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents