

#### www.immunoway.com.cn

# CD276 Rabbit pAb

CatalogNo: YT5285

## Key Features

Host Species <ul> <li>Rabbit</li> </ul>	<ul><li>Reactivity</li><li>Human,Mouse,Rat</li></ul>	Applications <ul> <li>WB,IHC,IF,ELISA</li> </ul>
MW • 100kD (Observed)	Isotype • IgG	

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC: 1:100-300 ELISA 1:20000 IF 1:50-200

### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

### Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD276. AA range:271-320
Specificity	CD276 Polyclonal Antibody detects endogenous levels of CD276 protein.

### **Target Information**

#### Gene name CD276

Protein Name CD276 antigen

Organism	Gene ID	UniProt ID		
Human	<u>80381;</u>	<u>Q5ZPR3;</u>		
Mouse	<u>102657;</u>	<u>Q8VE98;</u>		
Rat	<u>315716;</u>	<u>Q7TPB4;</u>		

#### Cellular Membrane ; Single-pass type I membrane protein .

#### Localization

- **Tissue specificity** Ubiquitous but not detectable in peripheral blood lymphocytes or granulocytes. Weakly expressed in resting monocytes. Expressed in dendritic cells derived from monocytes. Expressed in epithelial cells of sinonasal tissue. Expressed in extravillous trophoblast cells and Hofbauer cells of the first trimester placenta and term placenta.
- **Function** Function: May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling., induction: By LPS in monocytes and by ionomycin in T and B lymphocytes. Up-regulated in cells mediating rejection of human transplants...miscellaneous:B7-H3 locus underwent genomic duplication leading to tandemly repeated immunoglobulin-like V and C domains (VC domains). The dominantly expressed human B7-H3 isoform contains tandemly duplicated VC domains. In contrast, mouse B7-H3 transcript contains only one single VC domain form due to an exon structure corresponding to V domain-(pseudoexon C)-(pseudoexon V)-C domain. This duplication appearing in primates is suggested to be very recent supporting a model of multiple independent emergence of tandem VC repeats within human and monkey species., similarity: Belongs to the immunoglobulin superfamily. BTN/MOG family., similarity: Contains 2 Ig-like C2-type (immunoglobulin-like) domains., similarity: Contains 2 Ig-like V-type (immunoglobulin-like) domains., subunit: Interacts with TREML2 and this interaction enhances T-cell activation.,tissue specificity:Ubiquitous but not detectable in peripheral blood lymphocytes or granulocytes. Weakly expressed in resting monocytes. Expressed in dendritic cells derived from monocytes. Expressed in epithelial cells of sinonasal tissue. Expressed in extravillous trophoblast cells and Hofbauer cells of the first trimester placenta and term placenta..

### Validation Data



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100

	293
(kD)	
117-	11111
85-	
48-	
24	
34-	
26-	
20-	
19-	

Western blot analysis of lysate from 293 cells, using CD276 Antibody.

### **Contact information**

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **CD276 Rabbit pAb**  For Research Use Only. Not for Use in Diagnostic Procedures.

Immunoway - 4 / 4