

QKI Rabbit pAb

CatalogNo: YT7323

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA, IHC

MW

- 38kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human QKI AA range: 198-248

Specificity This antibody detects endogenous levels of QKI at Human/Mouse/Rat

Target Information

Gene name QKI HKQ

Protein Name QKI


Organism	Gene ID	UniProt ID
Human	9444;	Q96PU8;
Mouse	19317;	Q9QYS9;
Rat		Q91XU1;

Cellular Localization Nucleus . Cytoplasm .


Tissue specificity Expressed in the frontal cortex of brain. Down-regulated in the brain of schizophrenic patients.

Function Disease:Deletion of the QKI gene may play a role in astrocytic tumors.,Disease:May play a role in genetic susceptibility to schizophrenia. QKI levels of some isoforms such as isoform 6, are frequently down-regulated in schizophrenic patients.,Function:RNA-binding protein that plays a central role in myelination. Binds to the 5'-NACUAAAY-N(1,20)-UAAAY-3' RNA core sequence. Acts by regulating pre-mRNA splicing, mRNA export, mRNA stability and protein translation. Required to protect and promote stability of mRNAs such as MBP and CDKN1B which promotes oligodendrocyte differentiation. Participates in mRNA transport by regulating the nuclear export of MBP mRNA. Also involved in regulation of mRNA splicing of MAG pre-mRNA. Acts as a translational repressor.,PTM:Methylated by PRMT1.,PTM:Tyrosine phosphorylated at its C-terminus, probably by FYN. Phosphorylation leads to decreased mRNA-binding affinity, affecting transport and/or stabilization of MBP mRNA.,sequence Caution:Chimeric cDNA.,similarity:Contains 1 KH domain.,subunit:Homodimer. Does not require RNA to homodimerize. Able to heterodimerize with BICC1.,tissue specificity:Expressed in the frontal cortex of brain.,

Validation Data



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
QKI Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)