

InVivo Anti-Mouse PDL1 Antibody (10F.9G2)

CatalogNo: YV0027

| Key Features

Host Species

- Mouse

Reactivity

- Mouse

Applications

- In vivo

Isotype

- IgG2a, kappa

| Storage

Storage*

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from the date of receipt.

Formulation

0.01M PBS, pH 7.4.

| Recommended Dilution Ratios

| Basic Information

Purity

Endotoxin level <1 EU/mg. Purity>95% as determined by SDS-PAGE.

Clonality

Monoclonal

| Immunogen Information

| Target Information

Gene name

CD274 B7H1 PDCD1L1 PDCD1LG1 PDL1

Protein Name	Programmed cell death 1 ligand 1 (PD-L1) (PDCD1 ligand 1) (Programmed death ligand 1) (B7 homolog 1) (B7-H1) (CD antigen CD274)		
	Organism	Gene ID	UniProt ID
	Human	29126 ;	Q9NZQ7 ;
Cellular Localization	Cell membrane ; Single-pass type I membrane protein . Early endosome membrane ; Single-pass type I membrane protein . Recycling endosome membrane ; Single-pass type I membrane protein . Associates with CMTM6 at recycling endosomes , where it is protected from being targeted for lysosomal degradation. . ; [Isoform 1]: Cell membrane ; Single-pass type I membrane protein . ; [Isoform 2]: Endomembrane system ; Single-pass type I membrane protein .		
Tissue specificity	Highly expressed in the heart , skeletal muscle , placenta and lung. Weakly expressed in the thymus , spleen , kidney and liver. Expressed on activated T- and B-cells , dendritic cells , keratinocytes and monocytes.		
Function	Function:Involved in the costimulatory signal , essential for T-cell proliferation and production of IL10 and IFNG , in an IL2-dependent and a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation and cytokine production. ,induction:Up-regulated on T- and B-cells , dendritic cells , keratinocytes and monocytes after LPS and IFNG activation. Up-regulated in B-cells activated by surface Ig cross-linking. ,similarity:Belongs to the immunoglobulin superfamily. BTN/MOG family. ,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain. ,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain. ,subunit:Interacts with PDCD1. ,tissue specificity:Highly expressed in the heart , skeletal muscle , placenta and lung. Weakly expressed in the thymus , spleen , kidney and liver. Expressed on activated T- and B-cells , dendritic cells , keratinocytes and monocytes. ,		

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
InVivo Anti-Mouse PDL1 Antibody (10F.9G2)