

# Cyclophilin B (2B10) Mouse mAb (AbFluor 594)

CatalogNo: YM2020

#### **Key Features**

**Host Species** 

Mouse

Reactivity

Human, Rat, Mouse

Applications
• IF,WB,IHC,

IsotypeIgG

ConjugateAbFluor 594

## **Recommended Dilution Ratios**

Optimal working dilutions should be determined experimentally by the investigator Suggested starting dilutions are as follows:IHC 1:50-300 IF 1:50-200.

## Storage

Storage\*

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Store in dark.

**Formulation** 

Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.

## **Basic Information**

**Clonality** 

Monoclonal

**Clone Number** 

2B10

## Immunogen Information

**Specificity** 

Cyclophilin B Monoclonal Antibody(2B10) AbFluor™ 594 Conjugated specially designed for your Immunofluorescence analysis.

## | Target Information

Gene name

**PPIB** 

**Protein Name** 

Peptidyl-prolyl cis-trans isomerase B (PPlase B) (CYP-S1) (Cyclophilin B) (Rotamase B) (S-

cyclophilin) (SCYLP)

Organism Gene ID UniProt ID

Human 5479; P23284;

Cellular Localization Virion . (Microbial infection).; Endoplasmic reticulum lumen . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). .

**Tissue specificity** Brain, Fetal brain cortex, Prostate, Skin,

**Function** Catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,Caution:It is

uncertain whether Met-1 or Met-9 is the initiator.,enzyme regulation:Cyclosporin A (CsA) inhibits CYPB.,Function:PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,similarity:Belongs to the cyclophilin-type PPlase family. PPlase B subfamily.,similarity:Contains 1 PPlase cyclophilin-type domain.,subcellular location:Identified by mass spectrometry in melanosome fractions

from stage I to stage IV.,

#### Validation Data

#### | Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Cyclophilin B (2B10) Mouse mAb (AbFluor 594)

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents