

## $\beta$ -Tubulin (5G3) Mouse mAb (AbFluor 555)

CatalogNo: YM2199

### | Key Features

**Host Species**

- Mouse

**Reactivity**

- Human, Rat, Mouse, Mk, Dg, Ch, Hamster, Rabbit, sheep, Insect, Yeast

**Applications**

- WB, IF, IHC

**Isotype**

- IgG1

**Conjugate**

- AbFluor 555

### | Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator

Suggested starting dilutions are as follows: IHC 1:200

IF 1:200.

### | Storage

**Storage\***

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Store in dark.

**Formulation**

Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.

### | Basic Information

**Clonality**

Monoclonal

**Clone Number**

5G3

### | Immunogen Information


**Specificity**

$\beta$ -Tubulin Monoclonal Antibody(5G3) AbFluor™ 555 Conjugated specially designed for your Immunofluorescence analysis.

## Target Information

Gene name	TUBB3		
Protein Name	Tubulin beta-3 chain		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">10381</a> ;	<a href="#">Q13509</a> ;
Cellular Localization	Cytoplasm, cytoskeleton . Cell projection, growth cone . Cell projection, lamellipodium . Cell projection, filopodium .		
Tissue specificity	Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.		
Function	Domain:The highly acidic C-terminal region may bind cations such as calcium.,Function:Receptor for MSH (alpha, beta and gamma) and ACTH. The activity of this receptor is mediated by G proteins which activate adenylate cyclase.,Function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,polymorphism:Genetic variations in MC1R are associated with variation in skin/hair/eye pigmentation type 2 (SHEP2) [MIM:266300]. Hair, eye and skin pigmentation are among the most visible examples of human phenotypic variation, with a broad normal range that is subject to substantial geographic stratification. In the case of skin, individuals tend to have lighter pigmentation with increasing distance from the equator. By contrast, the majority of variation in human eye and hair color is found among individuals of European ancestry, with most other human populations fixed for brown eyes and black hair.,polymorphism:Variations in MC1R are linked to the degree of skin pigmentation (Types I-IV). Type I skin the most lightly pigmented and type IV the most dark pigmented. Partial loss-of-function mutations are associated with fair skin, poor tanning and increased skin cancer risk.,similarity:Belongs to the G-protein coupled receptor 1 family.,similarity:Belongs to the tubulin family.,subunit:Dimer of alpha and beta chains.,tissue specificity:Melanocytes and corticoadrenal tissue.,		

## Validation Data



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**β-Tubulin (5G3)**  
**Mouse mAb**  
**(AbFluor 555)**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)