

Iba 1 (PTR1347) Mouse mAb

CatalogNo: YM4765

Key Features

Host Species

Mouse

MW

16kD (Calculated)
 17kD (Observed)

Reactivity

· Human, Mouse, Rat

Isotype

IgG1,Kappa

Applications

• IHC,WB,IF,ELISA

Recommended Dilution Ratios

IHC 1:200-1000 WB 1:500-2000 IF 1:100-500

ELISA 1:1000-5000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Basic Information

Clonality Monoclonal

Clone Number PTR1347

Immunogen Information

Immunogen Synthesized peptide derived from part region of human protein AA range: 50-147

Specificity This antibody detects endogenous levels of Iba 1 protein.

| Target Information

Gene name

AIF1 G1 IBA1

Protein Name

Allograft inflammatory factor 1 (AIF-1) (Ionized calcium-binding adapter molecule 1) (Protein

Organism	Gene ID	UniProt ID
Human	<u>199;</u>	<u>P55008;</u>
Mouse	<u>11629;</u>	<u>070200;</u>
Rat	29427;	<u>P55009;</u>

Cellular Localization


Membranous

Tissue specificity Detected in T-lymphocytes and peripheral blood mononuclear cells.


Function

Function: May play a role in macrophage activation and function., PTM: Phosphorylated on serine residues., similarity: Contains 2 EF-hand domains.,


Validation Data




Various whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-lba 1 (PTR1347) antibody. The HRPconjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: THP-1 Lane 2: mouse liver Lane 3: rat brain



Human tonsil tissue was stained with Anti-Iba 1 (PTR1347) Antibody



Human brain tissue was stained with Anti-Iba 1 (PTR1347) Antibody




Human kindey tissue was stained with Anti-Iba 1 (PTR1347) Antibody




Mouse brain tissue was stained with Anti-Iba 1 (PTR1347) Antibody




Rat brain tissue was stained with Anti-Iba 1 (PTR1347) Antibody




Human spleen tissue was stained with Anti-Iba 1 (PTR1347) Antibody



Human stomach tissue was stained with Anti-Iba 1 (PTR1347) Antibody



Fluorescence multiplex immunohistochemical analysis of Mouse brain tissue (formalin-fixed paraffin-embedded section). The immunostaining was performed by Sextuple-Fluorescence kit (RS0039, Immunoway). GFAP mouse mAb(YM426 Immunoway) green, S100 mouse mAb(YM6987 Immunoway) cyan,Neurofilament mouse mAb(YM6897 Immunoway) purple,Iba 1 mouse mAb(YM4765 Immunoway) red,CD8 a mouse mAb(YM4815 Immunoway) yellow, The section was incubated in 5 rounds of staining; sequentially for Anti-antibodies; each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Immunoway YS0004, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain. Microscopy and pseudocoloring of individual dyes was performed using a Slideviewer Imaging System (Excilone).



Fluorescence multiplex immunohistochemical analysis of Human tonsil tissue (formalin-fixed paraffin-embedded section). The immunostaining was performed by Pentuple-Fluorescence kit (RS0038, Immunoway). CK-pan mouse mAb(YM6815 Immunoway) green, Ki-67 rabbit mAb(YM7002 Immunoway) red, Iba 1 mouse mAb(YM4765 Immunoway) purple,CD8 a mouse mAb(YM4815 Immunoway) cyan, CD20 mouse mAb(YM4814 Immunoway) yellow, The section was incubated in 5 rounds of staining; sequentially for Anti-antibodies; each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Immunoway YS0004, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain. Microscopy and pseudocoloring of individual dyes was performed using a Slideviewer Imaging System (Excilone).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Iba 1 (PTR1347)

Mouse mAb