

TCF-4 (PT1740R) PT™ Rabbit mAb

CatalogNo: YM9582 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 71kD (Calculated)
- 62kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000**WB 1:2000-1:10000****IF 1:200-1:1000****ELISA 1:5000-1:20000**

Basic Information

Clonality Monoclonal**Clone Number** PT1740R

Immunogen Information

Specificity Endogenous

Target Information

Gene name TCF4

Protein Name Transcription factor 4


Organism	Gene ID	UniProt ID
Human	6925 ;	P15884 ;
Mouse	21413 ;	
Rat	84382 ;	Q62655 ;

Cellular Localization Nucleus .

Tissue specificity Expressed in adult heart, brain, placenta, skeletal muscle and to a lesser extent in the lung. In developing embryonic tissues, expression mostly occurs in the brain.

Function Disease:Defects in TCF4 are a cause of Pitt-Hopkins syndrome (PTHS) [MIM:610954].,Disease:Haploinsufficiency of TCF4 is a cause of Pitt-Hopkins syndrome (PTHS) [MIM:610954]. PTHS is a rare syndromic encephalopathy characterized by severe psychomotor delay, epilepsy, daily bouts of diurnal hyperventilation starting in infancy, mild postnatal growth retardation, postnatal microcephaly, and distinctive facial features. Since most hitherto reported cases have been sporadic, with males and females equally affected, PTHS is regarded as an autosomal dominant condition.,Function:Transcription factor that binds to the immunoglobulin enhancer Mu-E5/KE5-motif. Binds to the E-box present in the somatostatin receptor 2 initiator element (SSTR2-INR) to activate transcription (By similarity). Preferentially binds to either 5'-ACANNTGT-3' or 5'-CCANNTGG-3'.,sequence Caution:Incomplete and probable erroneous sequence.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Forms homo- or heterooligomers with myogenin. Interacts with HIVEP2.,tissue specificity:Expressed in adult heart, brain, placenta, skeletal muscle and to a lesser extent in the lung. In developing embryonic tissues, expression mostly occurs in the brain.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-TCF-4 (PT1740R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: CACO-2 Lane 2: Mouse brain Predicted band size: 71kDa Observed band size: 62kDa

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
TCF-4 (PT1740R)
PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)